

Suitability of A_4 as a Family Symmetry in Grand Unification

Ernest Ma

Physics Department, University of California, Riverside, California 92521

Abstract

In the recent successful applications of the non-Abelian discrete symmetry A_4 to the tribimaximal mixing of neutrinos, lepton doublets and singlets do not transform in the same way. It appears thus to be unsuitable as a family symmetry in grand unification. A simple resolution of this dilemma is proposed.

To discover a possible underlying symmetry for the neutrino mass matrix, the problem of having a charged-lepton mass matrix with three very different eigenvalues must be faced. A proven solution is to make use of a non-Abelian discrete symmetry, such as A_4 [1, 2]. This is the group of even permutations of four objects, which is also the symmetry group of the perfect tetrahedron [3]. It has 12 elements and 4 irreducible representations: $\mathbf{1}$, $\mathbf{1}'$, $\mathbf{1}''$, and $\mathbf{3}$, with the multiplication rule

$$\mathbf{3} \times \mathbf{3} = \mathbf{1} + \mathbf{1}' + \mathbf{1}'' + \mathbf{3} + \mathbf{3}. \quad (1)$$

Specifically, for $a_{1,2,3} \sim \mathbf{3}$, $b_{1,2,3} \sim \mathbf{3}$, and $\omega = \exp(2\pi i/3) = -1/2 + i\sqrt{3}/2$,

$$a_1 b_1 + a_2 b_2 + a_3 b_3 \sim \mathbf{1}, \quad (2)$$

$$a_1 b_1 + \omega^2 a_2 b_2 + \omega a_3 b_3 \sim \mathbf{1}', \quad (3)$$

$$a_1 b_1 + \omega a_2 b_2 + \omega^2 a_3 b_3 \sim \mathbf{1}'', \quad (4)$$

$$(a_2 b_3, a_3 b_1, a_1 b_2) \sim \mathbf{3}, \quad (5)$$

$$(a_3 b_2, a_1 b_3, a_2 b_1) \sim \mathbf{3}. \quad (6)$$

Let

$$(\nu_i, l_i) \sim \mathbf{3}, \quad l_i^c \sim \mathbf{1}, \mathbf{1}', \mathbf{1}'', \quad (\phi_i^0, \phi_i^-) \sim \mathbf{3}, \quad (7)$$

then the charged-lepton mass matrix is given by

$$\begin{aligned} \mathcal{M}_l &= \begin{pmatrix} f_1 v_1 & f_2 v_1 & f_3 v_1 \\ f_1 v_2 & f_2 \omega v_2 & f_3 \omega^2 v_2 \\ f_1 v_3 & f_2 \omega^2 v_3 & f_3 \omega v_3 \end{pmatrix} \\ &= \begin{pmatrix} v_1 & 0 & 0 \\ 0 & v_2 & 0 \\ 0 & 0 & v_3 \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix} \begin{pmatrix} f_1 & 0 & 0 \\ 0 & f_2 & 0 \\ 0 & 0 & f_3 \end{pmatrix}, \end{aligned} \quad (8)$$

where $\langle \phi_i^0 \rangle = v_i$. If $v_1 = v_2 = v_3 = v$ (which breaks A_4 to the residual symmetry Z_3), then

$$\mathcal{M}_l = U_L \begin{pmatrix} f_1 & 0 & 0 \\ 0 & f_2 & 0 \\ 0 & 0 & f_3 \end{pmatrix} \sqrt{3}v, \quad (9)$$

where

$$U_L = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix}. \quad (10)$$

This solves the problem of having a diagonal $\mathcal{M}_l$ with three independent eigenvalues despite the presence of an underlying symmetry.

In the neutrino sector, assuming that $\mathcal{M}_\nu$ is Majorana and allowing neutrino masses to come from Higgs triplets, it has been shown [4] that the form

$$\mathcal{M}_\nu = \begin{pmatrix} a+2b & 0 & 0 \\ 0 & a-b & d \\ 0 & d & a-b \end{pmatrix} \quad (11)$$

leads to tribimaximal mixing [5] as follows. Rotating $\mathcal{M}_\nu$ to the basis $(\nu_e, \nu_\mu, \nu_\tau)$ with U_L of Eq. (10), one obtains

$$\begin{aligned} \mathcal{M}_\nu^{(e,\mu,\tau)} &= U_L^\dagger \mathcal{M}_\nu U_L^* \\ &= \begin{pmatrix} a + (2d/3) & b - (d/3) & b - (d/3) \\ b - (d/3) & b + (2d/3) & a - (d/3) \\ b - (d/3) & a - (d/3) & b + (2d/3) \end{pmatrix}, \end{aligned} \quad (12)$$

which is diagonalized exactly, resulting in

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} \sqrt{2/3} & \sqrt{1/3} & 0 \\ -\sqrt{1/6} & \sqrt{1/3} & -\sqrt{1/2} \\ -\sqrt{1/6} & \sqrt{1/3} & \sqrt{1/2} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}, \quad (13)$$

where

$$m_1 = a - b + d, \quad (14)$$

$$m_2 = a + 2b, \quad (15)$$

$$m_3 = -a + b + d. \quad (16)$$

The tribimaximal pattern of Harrison, Perkins, and Scott [5] is thus achieved. However, the neutrino masses $m_{1,2,3}$ remain arbitrary. Nevertheless, in the special cases of $b = 0$ [6]

or $a = 0$ [7], predictions relating m_{ν_e} (the kinematic ν_e mass), m_{ee} (the effective Majorana neutrino mass in neutrinoless double beta decay), and Δm_{atm}^2 (the mass-squared-difference measured in atmospheric neutrino oscillations) are obtained [7].

Since (ν_i, l_i) and l_i^c transform differently under A_4 , this application is not compatible with grand unification in general. (An exception [8] is conceivable in $SU(5)$, where (ν_i, l_i) belong to the $\mathbf{5}^*$ representation and l_i^c to the $\mathbf{10}$.) It is thus desirable to consider instead

$$(\nu_i, l_i), l_i^c \sim \mathbf{3}. \quad (17)$$

In that case, the natural choice [9, 10] of the Higgs sector is

$$(\phi_i^0, \phi_i^-) \sim \mathbf{1}, \mathbf{1}', \mathbf{1}'', \quad (18)$$

which exhibits an already diagonal $\mathcal{M}_l$. Although the neutrino mixing pattern may still be consistent with data, the predictivity of Eq. (11) is now lost.

The resolution of this apparent dilemma is actually very simple. Instead of the Higgs assignments of Eq. (18), let

$$(\phi_i^0, \phi_i^-) \sim \mathbf{1}, \mathbf{3}, \quad (19)$$

instead. Then

$$\mathcal{M}_l = \begin{pmatrix} h_0 v_0 & h_1 v_3 & h_2 v_2 \\ h_2 v_3 & h_0 v_0 & h_1 v_1 \\ h_1 v_2 & h_2 v_1 & h_0 v_0 \end{pmatrix}. \quad (20)$$

Again for $v_1 = v_2 = v_3 = v$, it is exactly diagonalized:

$$\mathcal{M}_l = U_L \begin{pmatrix} h_0 v_0 + (h_1 + h_2)v & 0 & 0 \\ 0 & h_0 v_0 + (h_1 \omega + h_2 \omega^2)v & 0 \\ 0 & 0 & h_0 v_0 + (h_1 \omega^2 + h_2 \omega)v \end{pmatrix} U_R^\dagger, \quad (21)$$

where $U_L = U_R$ is given again by Eq. (10). This means that Eq. (11) for $\mathcal{M}_\nu$ again predicts tribimaximal mixing as desired. The obstacle to having a grand-unified model within this context is removed.

In the quark sector, $\mathcal{M}_u$ and $\mathcal{M}_d$ are also of the form of Eq. (20), thus predicting $V_{CKM} = 1$ in the limit of $v_1 = v_2 = v_3 = v$ as noted before [2]. This is not such a bad first approximation and it may be possible to recover a realistic V_{CKM} from explicit soft A_4 breaking terms.

In the context of the discrete symmetry $\Delta(27)$ which has recently been proposed [11, 12], the analogous $\mathbf{3} \times \mathbf{3} \times \mathbf{3}$ decomposition has 3 invariants, resulting in the form

$$\mathcal{M}_l = \begin{pmatrix} h_0 v_1 & h_1 v_3 & h_2 v_2 \\ h_2 v_3 & h_0 v_2 & h_1 v_1 \\ h_1 v_2 & h_2 v_1 & h_0 v_3 \end{pmatrix}, \quad (22)$$

which has the same limit as Eq. (20) for $v_1 = v_2 = v_3 = v$. However, Eq. (11) for $\mathcal{M}_\nu$ is not as easy to obtain as in A_4 .

In conclusion, thanks to Eq. (21), it has been shown that A_4 is a suitable candidate family symmetry in grand unification, allowing all quarks and leptons to transform according to its $\mathbf{3}$ representation, and predicting the tribimaximal mixing of neutrinos.

This work was supported in part by the U. S. Department of Energy under Grant No. DE-FG03-94ER40837.

References

- [1] E. Ma and G. Rajasekaran, Phys. Rev. **D64**, 113012 (2001).
- [2] K. S. Babu, E. Ma, and J. W. F. Valle, Phys. Lett. **B552**, 207 (2003).
- [3] E. Ma, Mod. Phys. Lett. **A17**, 2361 (2002).
- [4] E. Ma, Phys. Rev. **D70**, 031901(R) (2004).
- [5] P. F. Harrison, D. H. Perkins, and W. G. Scott, Phys. Lett. **B530**, 167 (2002). See also X.-G. He and A. Zee, Phys. Lett. **B560**, 87 (2003).
- [6] G. Altarelli and F. Feruglio, Nucl. Phys. **B720**, 64 (2005); *ibid.* **B741**, 215 (2006).
- [7] E. Ma, Phys. Rev. **D72**, 037301 (2005).
- [8] E. Ma, H. Sawanaka, and M. Tanimoto, hep-ph/0606103.
- [9] M. Hirsch, E. Ma, A. Villanova del Moral, and J. W. F. Valle, Phys. Rev. **D72**, 091301(R) (2005).
- [10] E. Ma, Mod. Phys. Lett. **A20**, 2601 (2005).
- [11] I. de Medeiros Varzielas, S. F. King, and G. G. Ross, hep-ph/0607045.
- [12] E. Ma, hep-ph/0607056.